

2-Amino-4-(nitroalkyl)-4H-chromene-3-carbonitriles as New Cytotoxic Agents

Afsaneh Zonouzi^a, Roghieh Mirzazadeh^a, Maliheh Safavi^b,
Sussan Kabudanian Ardestani^c, Saeed Emami^d and Alireza Foroumadi^{b,e*}

^a School of Chemistry, University College of Science, University of Tehran, P.O. Box 14155-6455, Tehran, Iran. ^b Faculty of Pharmacy and Pharmaceutical Sciences Research Center, Tehran University of Medical Sciences, Tehran, Iran. ^c Institute of Biochemistry and Biophysics, Department of Biochemistry, University of Tehran, Tehran, Iran. ^d Department of Medicinal Chemistry and Pharmaceutical Sciences Research Center, Faculty of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran. ^e Drug Design and Development Research Center, Tehran University of Medical Sciences, Tehran, Iran.

Abstract

A series of 2-amino-4-(nitroalkyl)-4H-chromene-3-carbonitriles were synthesized by an efficient multicomponent reaction in aqueous media using DBU as a catalyst at room temperature. Mild condition, environment friendly procedure and excellent yields are the main advantages of this procedure. The cytotoxic activity of target compounds were evaluated against three cancer cell lines MDA-MB-231, MCF-7 and T47D in comparison with etoposide as reference drug. Generally, all compounds showed good cell growth inhibitory activity with IC₅₀ values less than 30 µg/mL. Their activities were comparable or more potent than standard drug etoposide. The 6-bromo- derivatives 7e and 7f showed promising cytotoxic activity with IC₅₀ values in the range of 3.46–18.76 µg/mL, being more potent than etoposide against all tested cell lines.

Keywords: 4H-chromenes; Benzopyran; DBU; One-pot synthesis; Cytotoxic activity.

Introduction

4H-Chromene derivatives are important scaffold in organic and medicinal chemistry. They belong to a class of naturally occurring benzopyran derivatives with a wide range of biological applications, such as antiallergic (1), anti-proliferative (2), anticancer (3, 4), antibacterial (5, 6), antiviral (7) and potent apoptosis inducers (8). Such diverse biological activities have made chromene derivatives important for further development in medicinal

and organic synthesis studies (9, 10). In particular, 2-amino-4H-chromenes are of recent interest for their cytotoxic activities (11, 12); other biological activities have been observed, for instance, pyranopyranone 1 that served as precursor for the blood anticoagulant warfarin (13), benzopyrane 2 has been known for anticancer therapeutic (14) and (4H-chromen-4-yl) cyanoacetate 3 as inhibitor of Bcl-2 protein and apoptosis inducer (Figure 1) (15).

Multicomponent reactions have been successfully employed to generate highly diverse combinatorial libraries for high-throughput screening of biological and pharmacological activities (16, 17). This type of reaction

* Corresponding author:

E-mail: aforoumadi@yahoo.com

becomes increasingly important in organic and medicinal chemistry because it allows obtaining highly sophisticated polyfunctional molecules through simple one-pot procedures (18, 19). Multicomponent reaction protocol with environmentally benign solvents and catalytic systems is one of the most suitable strategies, which meets the requirements of green aspects of chemistry for developing libraries of medicinal scaffolds. Developing organic reactions in water has become highly popular in recent years due to its specific properties in mediating organic reactions and its friendliness to the environment (20-22).

Several procedures for the multicomponent preparation of 2-amino-4*H*-chromenes have been described (23-25). Earlier Elinson et al. have reported synthesis of 2-amino-4*H*-chromenes in the presence of NaOAc or KF as base (26). This method has been directed to offer corresponding 4*H*-chromenes by using malononitrile or cyanoacetate as one of C-H acids and exchange of substitution on 4-position.

Thus, considering the fact that the discovery of a novel anticancer drug is a urgent need (27, 28), in continuation of our research program to find a novel anticancer agent (29, 30), we developed a general rapid, easy and environmentally benign synthetic protocol for the synthesis of functionalized chromenes bearing 4-nitroalkyl moiety instead of the 4-aryl ring of cytotoxic agents 2-amino-4-aryl-4*H*-chromene-3-carbonitriles.

Experimental

Chemistry

Chemicals and solvents were obtained from Merck (Germany) and Fluka (Switzerland) and were used without further purification. Column chromatography was performed on silica gel (0.015-0.04 mm, mesh-size) and TLC on precoated plastic sheets (25 DCUV-254). Melting points were measured on a Barnstead Electrothermal melting point apparatus and are not corrected. Elemental analyses for C, H and N were performed using a Thermo Finnigan Flash EA1112 instrument. IR spectra were recorded on a Shimadzu FT-IR-4300 spectrophotometer as

KBr discs. ¹H NMR spectra were determined in CDCl₃ on a Bruker 500 spectrophotometer and chemical shifts are expressed in ppm downfield from tetramethylsilane. Mass spectra were recorded on a Finnigan-MAT 8430 spectrometer at an ionization potential of 70 ev.

General procedure for the synthesis of 2-amino-4-(nitroalkyl)-4*H*-chromene-3-carbonitrile derivatives (7a-f)

To a magnetically stirred mixture of salicylaldehyde 4 (1 mmol), malononitrile 5 (1 mmol) and nitroalkane 6 (2 mmol) in water (5 mL), 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 30 mol%) was added. The reaction mixture was stirred for 6 h at room temperature. The mixture was extracted with EtOAc (6 mL) and the organic phase was dried over Na₂SO₄, concentrated and purified on a silica gel column using EtOAc/hexane (4:1) as eluent to afford the product 7.

2-Amino-4-(nitromethyl)-4*H*-chromene-3-carbonitrile (7a)

White powder; yield 85%; m.p. 138-139 °C (Lit. 139-140 °C) (25); ¹H NMR (500 MHz, CDCl₃): 4.29 (dd, *J* = 6.7 and 4.7 Hz, 1H), 4.50 (dd, *J* = 12.5 and 6.8 Hz, 1H), 4.60 (dd, *J* = 12.5 and 4.7 Hz, 1H), 5.57 (br s, 2H), 6.9 (d, *J* = 8.7 Hz, 1H), 7.29-7.30 (m, 2H), 7.40 (dd, *J* = 8.7 and 2.3 Hz, 1H).

2-Amino-4-(1-nitroethyl)-4*H*-chromene-3-carbonitrile (7b)

White powder; yield 80%; diastereomeric ratio 1.2:1; m.p. 165-166 °C; major diastereomer: ¹H NMR (500 MHz, CDCl₃): 1.60 (d, *J* = 6.5 Hz, 3H), 4.2 (d, *J* = 6.5 Hz, 1H), 4.54-4.56 (m, 1H), 4.92 (br s, 2H), 6.98-7.35 (m, 4H); minor diastereomer: ¹H NMR (500 MHz, CDCl₃): 1.37 (d, *J* = 6.5 Hz, 3H), 4.4 (d, *J* = 3.5 Hz, 1H), 4.71-4.73 (m, 1H), 4.88 (br s, 2H), 6.98-7.35 (m, 4H); IR (KBr): 3429, 3324 (NH₂), 3027, 2993 (CH), 2201 (CN), 1604, 1571 (C=C), 1558 and 1381 (NO₂); MS (m/z): 245 (M⁺), 218 (M⁺-HCN), 215 (M⁺-NO), 199 (M⁺-NO₂), 172 (M⁺-NO₂, HCN), 144 (M⁺-HCN, CH₃CHNO₂), 114 (C₉H₆⁺), 77, 57. Anal. Calcd (%) for C₁₃H₁₃N₃O₃: C, 58.77; H, 4.52; N, 17.13. Found: C, 58.72; H, 4.45; N, 17.17.

2-Amino-6-chloro-4-(nitromethyl)-4H-chromene-3-carbonitrile (7c)

Yellow powder; yield 83%; m.p. 181-182 °C; ¹H NMR (500 MHz, CDCl₃): 4.33 (t, 1H), 4.55 (dd, *J* = 12.5 and 6.5 Hz, 1H), 4.63 (dd, *J* = 12.8 and 4.0 Hz, 1H), 4.84 (br s, 2H), 7.0 (d, *J* = 8.5 Hz, 1H), 7.18 (d, *J* = 2.5 Hz, 1H), 7.28-7.30 (m, 2H), 7.40 (dd, *J* = 8.7 and 2.3 Hz, 1H).

2-Amino-6-chloro-4-(1-nitroethyl)-4H-chromene-3-carbonitrile (7d)

Yellow powder; yield 75%; diastereomeric ratio 1:1; m.p. 166-167 °C; ¹H NMR (500 MHz, CDCl₃): 1.42 (d, *J* = 7.0 Hz, 3H), 1.60 (d, *J* = 6.5 Hz, 1H), 4.2 (d, *J* = 6.5 Hz, 1H), 4.36 (d, *J* = 3.5 Hz, 1H), 4.54-4.56 (m, 1H), 4.70-4.72 (m, 1H), 4.88 (br s, 2H), 6.99-7.31 (m, 4H); MS (*m/z*): 279, 281 (M⁺, M⁺+2), 233, 235 (M⁺, M⁺+2-NO₂), 205, 207 (M⁺, M⁺+2-CH₃-CH-NO₂), 171, 179 (M⁺-HCN, CH₃CHNO₂), 114 (C₉H₆⁺), 77, 57. Anal. Calcd (%) for C₁₂H₁₀ClN₃O₃: C, 51.52; H, 3.60; N, 15.02. Found: C, 51.49; H, 3.64; N, 15.07.

2-Amino-6-bromo-4-(nitromethyl)-4H-chromene-3-carbonitrile (7e)

White powder; yield 85%; m.p. 192-193 °C; ¹H NMR (500 MHz, CDCl₃): 4.39 (t, 1H), 4.51 (dd, *J* = 13.75 and 7.0 Hz, 1H), 4.63 (dd, *J* = 12.5 and 4.5 Hz, 1H), 4.85 (br s, 2H), 7.0 (d, *J* = 8 Hz, 1H), 7.17-7.20 (m, 2H), 7.33 (t, 1H); IR (KBr): 3440, 3326 (NH₂), 3030, 2998 (CH), 2204 (CN), 1608, 1574 (C=C), 1572, 1377 (NO₂), 812.

2-Amino-6-bromo-4-(1-nitroethyl)-4H-chromene-3-carbonitrile (7f)

White powder; yield 75%; diastereomeric ratio 1.6:1; m.p. 178-179 °C; major diastereomer: ¹H NMR (500 MHz, CDCl₃): 1.60 (d, *J* = 7.0 Hz, 3H), 4.18 (d, *J* = 6.5 Hz, 1H), 4.54-4.56 (m, 1H), 4.89 (br s, 2H), 6.93-7.45 (m, 4H); minor diastereomer: ¹H NMR (500 MHz, CDCl₃): 1.43 (d, *J* = 6.5 Hz, 3H), 4.35 (d, *J* = 3.5 Hz, 1H), 4.70-4.72 (m, 1H), 4.86 (br s, 2H), 6.93-7.45 (m, 4H); IR (KBr): 3432, 3325 (NH₂), 3029, 2996 (CH), 2194 (CN), 1600 (C=C), 1542, 1386 (NO₂), 820; MS (*m/z*): 322, 324 (M⁺, M⁺+2), 263, 265 (M⁺, M⁺+2-NO, HCN), 249, 251 (M⁺-NO₂, HCN), 221, 223 (M⁺, M⁺+2-HCN, CH₃CHNO₂), 170 (M⁺-HBr, CH₃CHNO₂), 143 (M⁺-HCN, HBr,

CH₃CHNO₂), 114 (C₉H₆⁺), 77, 57. Anal. Calcd (%) for C₁₂H₁₀BrN₃O₃: C, 44.47; H, 3.11; N, 12.96. Found: C, 44.51; H, 3.08; N, 12.99.

*Biological activity**Cell lines and cell culture*

The cell lines were purchased from the National Cell Bank of Iran (NCBI). The cells were grown in RPMI-1640 medium (GibcoBRL, UK) supplemented with 10% heat-inactivated fetal calf serum (GibcoBRL, UK) and 100 mg/mL streptomycin and 100U/mL penicillin at 37 °C in a humidified atmosphere with 5% CO₂ in air.

In-vitro cytotoxicity assay

The in vitro cytotoxic activity of each synthesized compounds 7a-f were assessed in comparison with etoposide using MTT colorimetric assay, first described by Mosmann (31) with modifications. Briefly, cultures in the exponential growth phase were trypsinized and diluted in complete growth medium to give a total cell count of 5×10⁴ cells/mL. 195 μL of suspension was added to wells of sterile 96-well plates (NUNC, Denmark) and allowed to attach overnight. The stock solutions of compounds were prepared in DMSO and then serially diluted with growth medium. After plating, 5 μL of a serial dilution of every compound was added. Each compound dilution was assessed in triplicate. The maximum amount of DMSO in the cell culture was 1%. Etoposide was used as positive control for cytotoxicity while three wells containing tumor cells cultured in 200 μL of complete medium were used as controls for cell viability. The plates were then incubated for 48 h. After incubation, 200 μL of RPMI-1640 without phenol red containing a final concentration of 0.5 mg/mL MTT (Sigma-Aldrich, Steinheim, Germany) was added to each well and the plate was incubated for another 4 h. After incubation, the culture medium was replaced with 100 μL of DMSO. Then the absorbance of each well was measured by using a microplate reader at 492 nm wavelengths. For each compound, the concentration causing 50% cell growth inhibition (IC₅₀) compared with the control was calculated from concentration response curves by regression analysis.




Figure 1. Structures of some 2-amino-4H-chromenes with diverse biological activities.

Results and Discussion


Chemistry

We report in this paper, highly efficient one-pot synthesis of 2-amino-4-(nitroalkyl)-4H-chromene derivatives 7a-f by using 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) as a catalyst in water at room temperature. A variety of bases with different pKa have been used for some multicomponent reactions. DBU base (pK_a = 12) is a neutral organic base with high basicity and has been used in many organic transformation in recent years (32). It is a sterically hindered amidine base and especially useful where side reactions due to the inherent nucleophilicity of basic nitrogen are a problem (33, 34).

In our attempts to develop an efficient

protocol, we focused on the efficient condensation of salicylaldehyde 4a, malononitrile 5 and nitromethane 6a in water at room temperature by several amounts of DBU as base (Figure 2). The best 85% yield of 2-amino-4-(nitromethyl)-4H-chromene-3-carbonitrile 7a was obtained when the reaction was performed using 30 mol% DBU at room temperature for 6 h. Under this optimal condition, salicylaldehydes 4, malononitrile 5 and nitroalkane 6 were condensed into corresponding substituted 2-amino-4-(nitroalkyl)-4H-chromenes 7a-f in 75-85% yields. The attempts with 3-, 4- or 5-methoxy derivatives of salicylaldehydes indicated that these aldehydes are not good candidates for this reaction. The 4-(nitroethyl)-products 7b, 7d and 7f were eluted from the column chromatography

Table 1. Cytotoxic activity (IC₅₀, in µg/mL)^a of compounds 7a-f against three cancer human cell lines in comparison with etoposide.



Compounds	X	R	MDA-MB-231	MCF-7	T47D
7a	H	H	19.47 ± 4.31	17.65 ± 2.34	21.75 ± 6.50
7b	H	CH ₃	20.38 ± 0.87	19.87 ± 4.31	29.12 ± 8.99
7c	Cl	H	9.44 ± 6.76	11.35 ± 0.77	13.56 ± 5.03
7d	Cl	CH ₃	11.74 ± 6.40	7.52 ± 4.44	12.05 ± 5.98
7e	Br	H	3.92 ± 0.96	7.08 ± 3.46	13.53 ± 7.83
7f	Br	CH ₃	3.46 ± 1.29	9.65 ± 2.26	18.76 ± 6.27
Etoposide	X	R	20.31 ± 2.14	18.43 ± 1.55	21.04 ± 2.55

^a The IC₅₀ values represent an average of three independent experiments (mean ± SD).




Figure 3. The proposed mechanism of DBU-catalyzed synthesis of 2-amino-4-(nitroalkyl)-4H-chromene-3-carbonitriles.

Conclusion

In conclusion, we have developed a facile, convenient and environmentally benign one-pot synthesis of multi-substituted chromenes using DBU as a catalyst in aqueous media at room temperature. In this instance, it is possible to apply the tenets of green chemistry to a medicinal setting in the development of new methodology to the biologically interesting molecules. Furthermore this strategy involves additional advantage over previous methods for proceeding under easy and mild conditions. All synthesised compounds showed good cytotoxic activity against MDA-MB-231, MCF-7

and T47D cell lines. Their activities were comparable or more potent than standard drug etoposide. These results indicate that 2-amino-4H-chromene-3-carbonitriles containing 4-(nitroalkyl)- moiety could serve as a new basis for the development of novel group of anticancer agents.

Acknowledgement

The support of this study by the Research Council of the University of Tehran is gratefully acknowledged.

References

- (1) Witte EC, Neubert P and Roesch A. 7-(Piperazinylpropoxy)-2H-1-benzopyran-2-ones. Ger. Offen. DE3427985, 1986. *Chem. Abstr.* (1986) 104: 224915f.
- (2) CP Dell and Smith CW. Antiproliferative derivatives of 4H-naphtho[1,2-b]pyran and process for their preparation, Eur. Pat. 537,94,9 21 Apr, 1993. *Chem. Abstr.* (1993) 119: 139102d.
- (3) Doshi JM, Tian D and Xing C. Structure-activity relationship studies of ethyl 2-amino-6-bromo-4-(1-cyano-2-ethoxy-2-oxoethyl)-4H-chromene-3-carboxylate (HA 14-1), an antagonist for antiapoptotic Bcl-2 proteins to overcome drug resistance in cancer. *J. Med. Chem.* (2006) 49: 7731-7739.
- (4) Kemnitzer W, Drewe J, Jiang S, Zhang H, Wang Y, Zhao J, Crogan-Grundy C, Xu L, Lamothe S, Gourdeau H, Denis R, Tseng B, Kasibhatla S and Cai SX. Discovery of 4-aryl-4H-chromenes as a new series of apoptosis inducers using a cell- and caspase-based high-throughput screening assay. 3. structure-activity relationships of fused rings at the 7,8-positions. *J. Med. Chem.* (2007) 50: 2858-2864.
- (5) Kumar RR, Perumal S, Senthilkumar P, Yogeewari P and Sriram D. An atom efficient, solvent-free, green synthesis and antimycobacterial evaluation of 2-amino-6-methyl-4-aryl-8-[(E)-arylmethylidene]-5,6,7,8-tetrahydro-4H-pyrano[3,2-c]pyridine-3-carbonitriles. *Bioorg. Med. Chem. Lett.* (2007) 17: 6459-6462.
- (6) Kidwai M, Saxena S, Khan MKR and Thukral SS. Aqua mediated synthesis of substituted 2-amino-4H-chromenes and *in-vitro* study as antibacterial agents. *Bioorg. Med. Chem. Lett.* (2005) 15: 4295-4298.
- (7) Martínez-Grau A and Marco JL. Friedländer reaction on 2-amino-3-cyano-4H-pyrans: Synthesis of derivatives of 4H-pyran[2,3-b]quinoline, new tacrine analogues. *Bioorg. Med. Chem. Lett.* (1997) 7: 3165-3170.
- (8) Gao M, Miller KD, Hutchins GD and Zheng Q-H. Synthesis of carbon-11-labeled 4-aryl-4H-chromens as new PET agents. *Appl. Radiat. Isot.* (2010) 68: 110-116.
- (9) Thompson LA. Recent applications of polymer-

- supported reagents and scavengers in combinatorial, parallel, or multistep synthesis. *Curr. Opin. Chem. Biol.* (2000) 4: 324-337.
- (10) Nefzi A, Ostresh JM and Houghten RA. The current status of heterocyclic combinatorial libraries. *Chem. Rev.* (1997) 97: 449-472.
- (11) Akbarzadeh T, Rafinejad A, Malekian Mollaghasem J, Safavi M, Fallah-Tafti A, Kabudanian Ardestani S, Shafiee A and Foroumadi A. 2-Amino-3-cyano-4-(5-arylisoxazol-3-yl)-4H-chromenes: Synthesis and in vitro cytotoxic activity. *Arch. Pharm.* (2012) 345: 386-392.
- (12) Rafinejad A, Fallah-Tafti A, Tiwari R, Shirazi AN, Mandal D, Shafiee A, Parang K, Foroumadi A, Akbarzadeh T. 4-Aryl-4H-naphthopyrans derivatives: One-pot synthesis, evaluation of Src kinase inhibitory and anti-proliferative activities. *DARU J. Pharm. Sci.* (2012) 20:100.
- (13) Wiener C, Schroeder CH, West BD and Link KP. Studies on the 4-hydroxycoumarins. XVIII. 3-[α -(acetamidomethyl)benzyl]-4-hydroxycoumarin and related products. *J. Org. Chem.* (1962) 27: 3086-3088.
- (14) Kemnitzer W, Kasibhatla S, Jiang S, Zhang H, Zhao J, Jia S, Xu L, Crogan-Grundy C, Denis R, Barriault N, Vaillancourt L, Charron S, Dodd J, Attardo G, Labrecque D, Lamothe S, Gourdeau H, Tseng B, Drewe J and Cai SX. Discovery of 4-aryl-4H-chromenes as a new series of apoptosis inducers using a cell- and caspase-based high-throughput screening assay. 2. Structure-activity relationships of the 7- and 5-, 6-, 8-positions. *Bioorg. Med. Chem. Lett.* (2005) 15: 4745-4751.
- (15) Wang J-L, Liu D, Zhang Z-J, Shan S, Han X, Srinivasula SM, Croce CM, Alnemri ES, and Huang Z. Structure-based discovery of an organic compound that binds Bcl-2 protein and induces apoptosis of tumor cells. *Proc. Natl. Acad. Sci. USA* (2000) 97: 7124-7129.
- (16) Saeedi M, Mahdavi M, Foroumadi A, Shafiee A. Synthesis of novel fused 4,5-dihydro-1,2,3-triazolo[1,5-a][1,4]benzodiazepine derivatives via four-component Ugi-Smiles-type reaction. *Tetrahedron* (2013) 69: 3506-3510.
- (17) Hosseini-Zare MS, Mahdavi M, Saeedi M, Asadi M, Javanshir S, Shafiee A, Foroumadi A. Synthesis of 2,3-diaryl-5H-imidazo[2,1-a]isoindol-5-ones via the one-pot reaction of 1,2-diketones, 2-formylbenzoic acids, and ammonium acetate. *Tetrahedron Lett.* (2012) 53: 3448-3451.
- (18) Dömling A. Recent advances in isocyanide-based multicomponent chemistry. *Curr. Opin. Chem. Biol.* (2002) 6: 306-313.
- (19) Dömling A and Ugi I. Multicomponent reactions with isocyanides. *Angew. Chem., Int. Ed.* (2000) 39: 3168-3210.
- (20) Breslow R. Water as a solvent for chemical reactions. In: *Green Chemistry*, Anastas PT and Williamson TC (Eds.) Oxford Press, New York (1998) pp 225-232.
- (21) Grieco PA. *Organic Synthesis in Water*; Blacky academic & Professional: London, (1998) 82-141.
- (22) Li CJ. Organic reactions in aqueous media with a focus on C-C bond formations: A decade update. *Chem. Rev.* (2005) 105: 3095-3165.
- (23) Makarem S, Mohammadi AA and Fakhari AR. A multi-component electro-organic synthesis of 2-amino-4H-chromenes. *Tetrahedron Lett.* (2008) 49: 7194-7196.
- (24) Maggi R, Ballini R, Sartori G and Sartorio R. Basic alumina catalysed synthesis of substituted 2-amino-2-chromenes via three-component reaction. *Tetrahedron Lett.* (2004) 45: 2297-2299.
- (25) Ballini R, Bosica G, Conforti ML, Maggi R, Mazzacani A, Righi P and Sartori G. Three-component process for the synthesis of 2-amino-2-chromenes in aqueous media. *Tetrahedron* (2001) 57: 1395-1398.
- (26) Elinson MN, Ilovaisky AI, Valentina MM Pavel AB Belyakov PA and Alexander OC. Solvent-free cascade reaction: direct multicomponent assembling of 2-amino-4H-chromene scaffold from salicylaldehyde, malononitrile or cyanoacetate and nitroalkanes. *Tetrahedron* (2010) 66: 4043-4048.
- (27) Miri R, Firuzi O, Peymani P, Nazarian Z, Shafiee A. Synthesis and Cytotoxicity Study of New Cyclopenta [b] quinoline-1,8-dione Derivatives. *Iranian J. Pharm. Res.* (2011) 10: 489-496.
- (28) Miri R, Javidnia K, Amirghofran Z, Salimi SH, Sabetghadam Z, Meilia S, Mehdipour AR. Cytotoxic Effect of Some 1, 4-Dihydropyridine Derivatives Containing Nitroimidazole Moiety. *Iranian J. Pharm. Res.* (2011) 10: 497-503.
- (29) Vosooghi M, Yahyavi H, Divsalar K, Shamsa H, Kheirollahi A, Safavi M, Ardestani SK, Sadeghi-Neshat S, Mohammadhosseini N, Edraki N, Khoshneviszadeh M, Shafiee A, Foroumadi A. Synthesis and In vitro cytotoxic activity evaluation of (E)-16-(substituted benzylidene) derivatives of dehydroepiandrosterone. *DARU J. Pharm. Sci.* (2013) 21:34.
- (30) Noushini S, Emami S, Safavi M, Ardestani SK, Gohari AR, Shafiee A, Foroumadi A. Synthesis and cytotoxic properties of novel (E)-3-benzylidene-7-methoxychroman-4-one derivatives. *DARU J. Pharm. Sci.* (2013) 21:31.
- (31) Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods* (1983) 65: 55-63.
- (32) Khurana JM, Nand B, Saluja P. DBU: a highly efficient catalyst for one-pot synthesis of substituted 3,4-dihydropyrano[3,2-c]chromenes, dihydropyrano[4,3-b]pyranes, 2-amino-4H-benzo[h]chromenes and 2-amino-4H benzo[g]chromenes in aqueous medium. *Tetrahedron* (2010) 66: 5637-5641.
- (33) Aggarwal VK, Mereu A. Superior amine catalysts for the Baylis-Hillman reaction: the use of DBU and its implications. *Chem. Commun.* (1999) 2311-2312.
- (34) Reed R, Réau R, Dahan F, Bertrand G. DBU and DBN are strong nucleophiles: X-ray crystal structures of onio- and dionio-substituted phosphanes *Angew. Chem., Int. Ed. Engl.* (1993) 32: 399-401.